



# A new generation detector for supersymmetric particles search by direct detection : MACHe3

Emmanuel Moulin, D. Santos, G. Perrin, G. Duhamel, F. Naraghi, Yu.M. Bunkov, H. Godfrin

## ► To cite this version:

Emmanuel Moulin, D. Santos, G. Perrin, G. Duhamel, F. Naraghi, et al.. A new generation detector for supersymmetric particles search by direct detection : MACHe3. COSLAB 2002 : Cosmology in the Laboratory, Sep 2002, Krakow, Poland. 21 Transparents. in2p3-00020396

**HAL Id: in2p3-00020396**


**<https://hal.in2p3.fr/in2p3-00020396>**

Submitted on 30 Jan 2004

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**A new generation detector for  
supersymmetric particles search  
by direct detection :  
**MACHe3**, **MA**trix of **C**ells of  
superfluid **He**l<sup>ium</sup> **3****



**ISN-CRTBT collaboration:**

- **ISN** : D. SANTOS, G. PERRIN, G. DUHAMEL,  
F. NARAGHI, E. MOULIN
- **CRTBT** : Yu. M. BUNKOV, H. GODFRIN


**Emmanuel MOULIN**

Institut des Sciences Nucléaires, CNRS/IN2P3  
and Université Joseph Fourier, Grenoble-France

# Main topics :

- Cosmological evidence for cold non-baryonic dark matter
- WIMP candidate : the neutralino
- MACHe3 project :
  - ➔ detection principle
  - ➔ detection threshold
  - ➔ events
- Results :
  - ➔ neutrons spectrum
  - ➔ simulation of muons (with GEANT3.21)
  - ➔ muons spectrum
- Conclusions and prospects


# Evidence for non-baryonic dark matter :



$$\Omega_{\text{tot}} = \Omega_{\gamma} + \Omega_{\Lambda} + \Omega_{\text{M}},$$

$$\Omega_{\text{tot}} \sim 1 \text{ (CMB)}$$

$\Omega_{\Lambda}$



$$\Omega_{\gamma} \sim 5 \times 10^{-5}$$

negligible

$$\Omega_{\Lambda} \sim 0.7 \text{ (SNIa)}$$

$$\Omega_{\text{M}} \sim 0.3$$

(clusters of galaxies)

$$\Omega_{\text{B}} \sim 0.04 \text{ (BBN)} :$$

$$\Omega_{\text{B}} < \Omega_{\text{M}}$$

$$\Omega_{\text{M}} = \Omega_{\text{B}} + \Omega_{\text{HDM}} + \Omega_{\text{CDM}}$$

$\Rightarrow$  existence of **non-baryonic** dark matter :

- hot, “HDM” : neutrinos,  $\Omega_{\nu} \leq 0.09$

- cold, “CDM” : **WIMPs**

# WIMPs candidate for cold non-baryonic dark matter : the neutralino $\chi$

- **cold** non-baryonic dark matter is favoured (structures formation in the Universe)
- characteristics of **WIMPs** ( **W**eakly **I**nteractive **M**assive **P**articles ) :
  - masses : from **30 GeV/c<sup>2</sup>** up to few **TeV/c<sup>2</sup>**
  - weak cross section : **< 10<sup>-2</sup> pb**
  - neutral of charge and color
- SUSY  $\Rightarrow$  lightest supersymmetric particle (LSP) : the **neutralino  $\chi$**  (with R-parity conserved)
  - belongs to WIMPs' class
  - its relic density:  $\Omega_{\chi} = O(10^{-1})$

# MACHe3 project for neutralinos $\chi$ detection from galactic halo :


- Measure of energy deposited by elastic scattering of  $\chi$  on  $^3\text{He}$  target nucleus
- $^3\text{He}$  as sensitive medium :
  - superfluid  $^3\text{He}$  :  $T \sim 100 \mu\text{K}$
  - very low energy gap of quasiparticles :  
threshold of 1 keV  
 $\Rightarrow$  ability of detecting weak recoil energies
- Main contributions to the background :
  - **gammas, neutrons, muons**
  - protons,  $\alpha$  particles (negligible)
  - micro-vibrations
- Main interesting features :
  - **high purity  $^3\text{He}$**
  - **spin 1/2** (axial interaction)
  - **neutron capture process**
  - **low sensitivity to  $\gamma$**

# CRTBT experimental hall :



- cryostat
- 5 cm thick lead shield

## Bolometer cell :



- Operating mode : Lancaster type bolometer **damping** effect on the vibrating wire of the quasiparticles cloud produced by an incoming particle interacting inside the cell
- Measure : damping linked to the **frequency width** of the vibrating wire calibrated in energy
- **Thermal diffusion** through the hole

# $\chi$ event in the granular detector :

- 2 contributions to the elastic scattering on nucleon :
  - **scalar** interaction :  $\sim 10^{-6}$  pb
  - **axial interaction** :  $\sim 10^{-2}$  pb

- Elastic scattering on  ${}^3\text{He}$  nucleus :

$$E_{\text{recoil}}^{\text{max}} = 2 \frac{m_{{}^3\text{He}} M_\chi^2}{\left(m_{{}^3\text{He}} + M_\chi\right)^2} v^2 \cong 2 m_{{}^3\text{He}} v^2$$

$\Rightarrow \chi$  event defined by an energy deposit  $\leq 6 \text{ keV}$

- The whole energy is deposited in a **single** cell  
 $\Rightarrow$  systematic discrimination compared with other events

- Expected event rate :


$$R = 1400 \times \sigma \text{ (pb)} / M_\chi \text{ (GeV/c}^2\text{)} [\text{kg}^{-1} \text{ day}^{-1}]$$

With  $\sigma \sim 10^{-2}$  pb and  $M_\chi \sim 30 \text{ GeV/c}^2$


$$\Rightarrow R \sim 7 \times 10^{-2} \text{ kg}^{-1} \text{ day}^{-1}$$



# Raw data at 100 $\mu\text{K}$ :




- ▶ low frequency modulation
- ▶ micro-vibrations



- ▶ very low micro-vibrations

# Sensitivity of the cell :



Acquisition spectrum at 100  $\mu$ K, without source :

- 3 peaks of about 10 keV
- detection of structures of about 1 keV

⇒ very promising results and :

- improvement of acquisition system
- understanding of micro-vibrations

# Discrimination of the different contributions of background for WIMPs detection :

➤  **$\gamma$  rays** (natural radioactivity) :

$^{40}\text{K}$ ,  $^{214}\text{Bi}$ ,  $^{214}\text{Pb}$ ,  $^{220}\text{Ac}$  et  $^{222}\text{Rn}$

➔ Compton effect  $\gg$  photoelectric effect :

$$\sigma_{\text{com}} / \sigma_{\text{pho}} \sim 10 \quad (\text{at } 100 \text{ keV})$$

➤ **neutrons** (considered as ultimate noise) :

➔ neutron capture process by the target nucleus, enhanced after thermalization :

$$\sigma_{\text{cap}} / \sigma_{\text{ela}} \sim 10 \quad (\text{at } 1 \text{ keV})$$

➤ **cosmic muons** (energy  $\sim 2 \text{ GeV}$ ) :

➔ energy loss in  $^3\text{He}$  by ionization


$$dE/dx \sim 0.1 \text{ MeV/cm}$$

# Neutrons separation by capture process by $^3\text{He}$ nucleus :

- ▶ Neutron capture : **exoenergetic** reaction




- ▶ Experiments with Am/Be source
- ▶ Acquisition time : 4.6 h, à 100  $\mu\text{K}$




- ▶ Position of the peak : **650 keV**, width **20 keV**  
⇒ energy resolution of **3 %**  
rate : 0.5  $\text{min}^{-1}$
- ▶ Shift compared with the expectation value of 764 keV:
  - vortices creation (Kibble mechanism)
  - UV photons emission

# Muons simulation :

- Estimate of the expectation counting rate by the single cell prototype :  $0.36 \text{ min}^{-1}$
- Simulation of muons inside the cell with **GEANT3.21** :
  - energy :  $2 \text{ GeV}$
  - draw of the muons generator :




➔ Peak at  $67 \pm 2.5 \text{ keV}$



# Muons spectrum :

- Acquisition time 19 h, at 100  $\mu$ K
- Peak at  $45 \pm 5$  keV
- Energies  $< 20$  keV : numerical limitation of the data analysis procedure



- Counting rate (muons +  $\gamma$ ) similar to estimate  $\sim 0.36 \text{ min}^{-1}$  (due to the fact of a low rate of  $\gamma$  :  $\sim 0.01 \text{ min}^{-1}$ )
- **Shift** compared with the simulation :
  - ➔ mechanism of UV photons emission ?

# Conclusions :


- Two major contributions to background are experimentally shown :
  - clear separation of thermal neutrons
  - detection of cosmic muons
- Simulation of muons inside the detector with GEANT3.21
- Estimate of a low counting rate of  $\gamma$  compared with the muons one

# Prospects :

- Improvement of the analysis method
- A better understanding of **background caused by micro-vibrations** is necessary (shape of these peaks would be more symmetric)
- Wavelets treatment is investigated
- **Calibration at low energy** : electron conversion source of  $^{57}\text{Co}$  with **7, 14**, 115 and 129 keV lines is considered
- **Multicellular** prototype in order to use correlations among the cells to improve discrimination of events



# Complementarity of MACHe3 with existing projects :



- exclusion limits from Edelweiss, CDMS experiments as well as the DAMA region
- dotted lines indicate projected limits of CRESST and CDMS experiments
- light points present SUSY models giving an event rate for **MACHe3** higher than 0.01 day<sup>-1</sup>

# Natural radioactivity :

- ▶ Experiment with a **Germanium detector** :
  - ⇒ radioactive contamination:  $^{40}\text{K}$ ,  $^{214}\text{Bi}$ ,  $^{214}\text{Pb}$  et  $^{220}\text{Ac}$
  - ⇒ counting rate
- ▶ GEANT3.21 simulation with :
  - Germanium cell
  - Helium cell
  - ⇒ ratio of the number of the detected counts
- ▶ Sources  $\gamma$  :  $^{137}\text{Cs}$  and  $^{60}\text{Co}$ 
  - counting rate :
    - $^{137}\text{Cs}$  :  $0.03 \text{ s}^{-1}$
    - $^{60}\text{Co}$  :  $0.01 \text{ s}^{-1}$
  - in agreement with estimate
- ▶  $\gamma$  counting rate without source :
  - $0.2 \text{ min}^{-1}$
  - **with a lead shield** (5 cm) :  $0.01 \text{ min}^{-1}$
  - ⇒ **low sensivity** of the detector for  $\gamma$  of natural radioactivity compared with muons

# $\chi$ event rate in the MACHe3 detector :

➤ Expression of event rate ( 1<sup>st</sup> approximation) :

$$R = \sigma \langle v \rangle ( \rho^0 / M_\chi ) \times ( M_{\text{det}} / m_{3\text{He}} )$$

thus,

$$R = 1400 \times \sigma \text{ (pb)} / M_\chi \text{ (GeV/c}^2\text{)} [\text{kg}^{-1} \text{ day}^{-1}]$$

with :

$$\begin{aligned} \langle v \rangle &\sim 270 \text{ km s}^{-1} \\ \rho^0 &\sim 0.3 \text{ GeV/c}^2 \text{ cm}^{-3} \\ m_{3\text{He}} &= 2.81 \text{ GeV/c}^2 \end{aligned}$$

➤ For  $\sigma \sim 10^{-2} \text{ pb}$  and  $M_\chi \sim 30 \text{ GeV/c}^2$

➔  $R \sim 7 \times 10^{-2} \text{ kg}^{-1} \text{ day}^{-1}$

# Evaluation the $\chi$ relic density in the Universe :

➤  $\chi$  **equilibrium** density :  $n_{\chi}^{eq} = g_{\chi} \left( \frac{m_{\chi} T}{2\pi} \right)^{3/2} e^{-m_{\chi}/T}$

➤ **Freezeout equation** :  $\Gamma = n_{\chi} \langle \sigma v \rangle = H$

resolution with  $n_{\chi}^{eq} \Rightarrow T_{freezeout} \cong m_{\chi} / 20$

(logarithmic corrections are neglected)

➤ For a **radiative** universe :  $H(T) = 1.66 g^{1/2} \frac{T^2}{M_{Pl}}$

➤ Density per covolume is **constant**  
(adiabatic expansion) :  $\left( \frac{n_{\chi}}{s} \right)_0 = \left( \frac{n_{\chi}}{s} \right)_{freezeout}$

with  $s \cong 0.4 g T^3$  and  $s_0 / k \cong 3000 cm^{-3}$

$$n_{\chi} = \frac{3 \times 10^{-8}}{m_{\chi} M_{Pl} \langle \sigma v \rangle}$$


and :  $\Omega_{\chi}^0 h_0^2 = \frac{O(10^{-27} cm^3 s^{-1})}{\langle \sigma v \rangle}$

$\Rightarrow \Omega_{\chi} = O(10^{-1})$


# Data analysis procedure in three steps :

- systematic subtraction of the **low frequency modulation** (polynom of order 5)
- **deconvolution**
- integration in order to limit the “**numerical**” **noise** caused by deconvolution


Raw data




Data after deconvolution



Data after integration



Energy distribution



# Principle of data analysis :

- raw spectrum  $f(t)$  : 
$$f(t) = \int_{-\infty}^{+\infty} h(u) g(t-u) du = (h * g)$$

with  $g(t)$  response **function of the wire**  
 $h(t)$  Dirac comb

- Fourier transform of  $f(t)$  : 
$$F(\omega) = H(\omega) \times G(\omega)$$

- with inverse Fourier transform : 
$$h(t) = TF^{-1}\left(\frac{F(\omega)}{G(\omega)}\right)$$

- use of a **reference peak** for deconvolution :

- at equal temperature, peaks have the **same** shape:

- ▶ rising time : 1 s

- ▶ descending time (at half height) : 10 s  
        (at 100 $\mu$ K)

- descending time : function of the running  
**temperature**